



Flowering rush

Butomus umbellatus (Aka grassy rush, water gladiolus)

Provincial Designation:
Fisheries Act, Weed
Control Act



Alvin Mitchell, Salish Kootenai College

<http://www.vashsad.ua/>

Overview:

Flowering rush is a cattail-like perennial of freshwater wetlands. It is native to Africa, Asia and Europe¹ and was likely introduced to North America as an ornamental plant. It is the only member of the Butomaceae family and is able to reproduce both by seed and vegetatively (rhizomatous roots form bulbils which separate from the parent plant³). Flowering rush infestations can displace native vegetation and result in reduced water quality which may disrupt valuable fish and wildlife habitat. Dense stands in irrigation ditches can reduce water availability and in lakes can interfere with boat propellers and swimming.³ Plants flower summer to fall.¹ Flowers are hermaphroditic (contain both male & female organs) and are pollinated by bees, flies and butterflies.²

Habitat:

Flowering rush can grow on water margins or as a submerged plant with flexible leaves suspended in deeper water (3-6 m).³ It is widely tolerant of soil types (sandy to clay

and soil acidity, but does require wet soil and full sun.⁴ It is hardy to Zone 2 in Canada.²

Identification:

Flowering rush can be confused with sedges when not in bloom but is usually a much larger plant.³

Stems: Are erect and triangular near the base. Plants grow to 150 cm.¹

Leaves: Are green and sword-shaped,⁵ originate from base of plant,² and are triangular in cross-section, twisted toward the tip, and feel spongy when compressed.³

Flowers: Are 2-2.5 cm wide with 3 slightly greenish sepals and 3 petals.³ Twenty to fifty pink through white flowers are borne on umbrella shaped clusters³. Anthers are red.⁵ There are 9 stamens ranged in an inner whorl of 3 and outer whorl of 6.³

Prevention:

Simply do not grow it - talk to your local nursery about non-invasive alternatives. Flowering rush can spread by seed or root frag-

ments so care must be taken with attempts to remove existing plants. It has been observed to invade aquatic areas with existing vegetation more slowly³ - maintain existing stands and prevent disturbance.

Control:

Mechanical: Cutting below the water surface can suppress plants but will need to be repeated. Hand digging is feasible with small infestations but care must be taken to remove all parts of the plant - root fragments can drift with water movement and result in new infestations². All plant matter should be removed and disposed of in landfill-bound garbage.

Chemical: Diquat is registered for use on flowering rush in lakes, ponds, irrigation canals and slow moving streams. Herbicide applications near water bodies require specific applicator certification and permits from Alberta Environment. Always check product labels to ensure the herbicide is registered for use on the target plant in Canada by the Pest Management Regulatory Agency. Always read and follow label directions. Con-

continued next page

Flowering Rush (Continued)

sult your local Agricultural Fieldman or Certified Pesticide Dispenser for more information.

Biological: None researched to date.



Stems

Don Battiste



Mature Flowers

Don Battiste




Root Mass

Leslie J. Mehrhoff, University of Connecticut, bugwood.org



Infestation

Leslie J. Mehrhoff, University of Connecticut, bugwood.org



Seeds

Steve Hurst, USDA NRCS plants database, bugwood.org



Nicole Kimmel, ARD

REFERENCES

- 1 Flora of North America. *Butomus umbellatus* www.eFloras.org
- 2 Global Invasive Species Database. *Butomus umbellatus*. www.issg.org
- 3 Montana State University Extension. Flowering Rush. <http://msuextension.org/publications/AgandNaturalResources/EB0201.pdf>
- 4 USDA Forest Service. Weed of the Week, Flowering Rush. www.na.fs.fed.us/fhp/invasive_plants
- 5 Hitchcock and Cronquist. Flora of the Pacific Northwest. 1973. University of Washington Press. p 557.